

Recombinant human R-spondin 3 protein (Qk032)



Type: Stem cells

Available for purchase: Unit Size (µg): 25, 50, 100, 500, 1000

Buy online with secure credit card or purchase order.

[View this product and buy online](#)

Product Information

Human R-spondin 3 protein potentiates Wnt signaling in and has been shown to function in crypt regeneration in the intestine and control stem cell and progenitor cell behavior during kidney development. R-spondin 3 is used alongside R-spondin 1 in intestinal [organoid](#) culture systems.

Qk032 is a 17 kDa highly pure, bioactive domain of human R-spondin 3 comprising the two cysteine-rich furin-like domains, which are necessary and sufficient for Wnt signalling potentiation and are the essential domains for activity in stem cell and organoid culture, [animal origin-free](#) and carrier-protein free.

This protein is also available as GMP compliant [Cell Therapy Grade](#), to enquire email ryan.weber@matriq.com.

Alternative protein names

R-spondin-3, RSPO3, RSPO-3, Protein with TSP type-1 repeat, PWTSR, Roof plate-specific spondin-3, Thrombospondin type-1 domain-containing protein 2, THSD2, , R spondin 3, Qk32

Molecular weight

17 kDa

Protein Uniprot number

High purity bioactive domain of human R-spondin 3 protein (Uniprot: Q9BXY4)

Species reactivity

- human
- species similarity:
- mouse - 97%
- rat - 97%
- bovine - 98%
- porcine - 98%

Product Information

- >98%, by SDS-PAGE quantitative densitometry
- Expressed in *E. coli*
- Animal origin-free (AOF) and carrier protein-free
- Manufactured in our Cambridge, UK laboratories
- Lyophilized from acetonitrile, TFA

Reconstitution instructions

- Resuspend in 10 mM HCl (Reconstitution solution A) at >50 µg/ml

Featured applications

- Chemically defined media for organoid culture
- Intestinal organoid culture
- Tumor organoid culture


Further quality assays

- Mass spectrometry: single species with expected mass
- Recovery from stock vial: >95%
- Endotoxin: <0.05 EU/µg protein

Scientific Information

Bioactivity

R-spondin 3 activity was determined using the Wnt-responsive firefly luciferase reporter assay. HEK293T cells transfected with reporter TOP-FLASH were treated in triplicate with increasing concentrations of R-spondin 3 in the presence of Wnt-conditioned media (1:8 dilution). RSP1 enhances Wnt- β catenin signaling in HEK293T cells. Cells were grown overnight, and luciferase activity is measured and normalized. EC50 = 0.3 nM (5 ng/ml). Data from Qk032 #010.



Purity

R-spondin 3 bioactive domain migrates at 17 kDa in non-reducing (- β ME) conditions and upon reduction (+ β ME). Purified recombinant protein (7 μ g) was resolved using 15% w/v SDS-PAGE in reduced (+ β -mercaptoethanol, R) and non-reduced conditions (NR) and stained with Coomassie Brilliant Blue R250. Data from Qk032 lot #010.



Original product page: <https://ryan.calliope-alpha.ts.net/product/recombinant-human-rspondin-3-protein/>

PDF generated: 12 May 2026

Copyright © 2026 by Qkine Ltd. All rights reserved including graphics and images.