

## Recombinant human OSM protein (Qk049)



**Type:** Stem cells

**Available for purchase:** Unit Size (µg): 25, 50, 100, 500, 1000

Buy online with secure credit card or purchase order.

[View this product and buy online](#)

### Product Information

---

Human OSM protein (oncostatin M) is used in the differentiation of human pluripotent stem cells into hepatocyte-like cells. OSM also plays roles in osteogenesis and neurogenesis, and is an important regulator of the [hematopoietic](#) stem cell niche in the bone marrow.

Qkine human oncostatin M protein is high purity and [animal origin-free](#) (AOF) and carrier-protein free for reproducible results.

**This protein is also available as GMP compliant [Cell Therapy Grade](#), to enquire email [ryan.weber@matriq.com](mailto:ryan.weber@matriq.com).**

#### Alternative protein names

Oncostatin M, OSM, Qk49

#### Molecular weight

22 kDa

#### Protein Uniprot number

High purity human OSM protein (Uniprot: P13725)

#### Species reactivity

- human
- species similarity:

- mouse - 50%
- rat - 50%
- bovine - 60%
- porcine - 68%
- NB: although the sequence identity is low, human OSM has been shown to be active on mouse ES cells

### **Product Information**

- >98%, by SDS-PAGE quantitative densitometry
- Expressed in *E. coli*
- Animal origin-free (AOF) and carrier protein-free
- Manufactured in our Cambridge, UK laboratories
- Lyophilized from acetonitrile, TFA

### **Reconstitution instructions**

- Resuspend in sterile-filtered water at >50 µg/ml

### **Featured applications**

- Maturation of hepatoblasts into hepatocyte like cells


### **Further quality assays**

- Mass spectrometry: single species with expected mass
- Recovery from stock vial: >95%
- Endotoxin: <0.05 EU/µg protein

## Scientific Information


### Bioactivity

OSM activity was determined using the Promega serum response element luciferase reporter assay (\*) in transfected HEK293T cells. EC50 = 0.75 ng/ml (34 pM). Cells were treated in triplicate with a serial dilution of OSM for 6 hours. Firefly luciferase activity was measured and normalized to the control Renilla luciferase activity. Data from Qk049 lot #104336. [\\*Promega pGL4.33\[Luc2P/SRE/Hygro\] #E1340](#)



### Purity

OSM migrates as a single band at 20 kDa in non-reducing (NR) conditions and 22 kDa upon reduction (R). The slight band under the main band in the reduced sample is likely an artefact of the reduction protocol. No contaminating protein bands are visible. Purified recombinant protein (3 µg) was resolved using 15% w/v SDS-PAGE in reduced (+β-mercaptoethanol, R) and non-reduced (NR) conditions and stained with Coomassie Brilliant Blue R250. Data from Qk049 batch #104336.



**Original product page:** <https://ryan.calliope-alpha.ts.net/product/recombinant-human-osm-protein-qk049/>

**PDF generated:** 12 May 2026

Copyright © 2026 by Qkine Ltd. All rights reserved including graphics and images.