

## Recombinant human IGF-2 protein (Qk118)



**Type:** Stem cells

**Available for purchase:** Unit Size (µg): 25, 50, 100, 500, 1000

Buy online with secure credit card or purchase order.

[View this product and buy online](#)

### Product Information

---

Insulin-like Growth Factor 2 (IGF-2) is a growth factor that promotes cell proliferation, differentiation, and survival, particularly during fetal development. IGF-2 also regulates tissue growth and muscle regeneration. Recombinant IGF-2 stimulates proliferation and differentiation in various stem cell cultures.

Qkine human IGF-2 is a highly pure 7.5 kDa monomer, [animal origin-free](#) (AOF) and carrier-protein-free (CF). Qkine has optimized the IGF-2 manufacturing process to produce a highly bioactive protein with excellent lot-to-lot consistency for enhanced experimental reproducibility.

**This protein is also available as GMP compliant [Cell Therapy Grade](#), to enquire email [ryan.weber@matriqx.com](mailto:ryan.weber@matriqx.com).**

#### Alternative protein names

Insulin-like growth factor II, IGF2, IGF-II, Somatomedin-A, T3M-11-derived growth factor, IGF 2

#### Molecular weight

7.5 kDa

#### Protein Uniprot number

High purity human IGF-2 protein (Uniprot: P01344)

#### Species reactivity

- human
- species similarity:
- mouse - 91%
- rat - 94%
- bovine - 95%
- porcine - 98%

### **Product Information**

- >98%, by SDS-PAGE quantitative densitometry
- Expressed in *E. coli*
- Animal origin-free (AOF) and carrier protein-free
- Manufactured in our Cambridge, UK laboratories
- Lyophilized from acetonitrile, TFA

### **Reconstitution instructions**

- Resuspend in sterile-filtered water at >50 µg/ml

### **Featured applications**

- Cellular proliferation, migration and survival
- iPSC and primary cell differentiation to muscle
- Osteogenic differentiation


### **Further quality assays**

- Mass spectrometry: single species with expected mass
- Recovery from stock vial: >95%
- Endotoxin: <0.05 EU/µg protein

## Scientific Information


### Bioactivity

Recombinant IGF-2 activity was determined using an SRE reporter assay on transfected MCF-7 cells. Cells were treated in triplicate with a serial dilution of IGF-2 for 4 hours. Firefly activity was measured and normalized to the control Renilla luciferase activity. Data from Qk118 lot 204687. EC50 = 58 ng/ml (7.8 nM)



### Purity

Recombinant IGF-2 migrates at approximately 7.5 kDa in reduced (R) and non-reduced (NR) conditions. No contaminating protein bands are present. The purified recombinant protein (3 µg) was resolved using 18% w/v SDS-PAGE in reduced (+β-mercaptoethanol, R) and non-reduced (NR) conditions and stained with Coomassie Brilliant Blue R250. Data from Qk118 lot #204687.



**Original product page:** <https://ryan.calliope-alpha.ts.net/product/recombinant-human-igf-2-protein-qk118/>

**PDF generated:** 12 May 2026

Copyright © 2026 by Qkine Ltd. All rights reserved including graphics and images.