

Recombinant mouse/rat FGF-8a protein (Qk020)



Type: Stem cells

Available for purchase: Unit Size (µg): 25, 50, 100, 500, 1000

Buy online with secure credit card or purchase order.

[View this product and buy online](#)

Product Information

Fibroblast growth factor 8a (FGF-8a) is a member of the [FGF family](#) and plays a key role in the regulation of embryogenesis, cellular proliferation, differentiation, and migration. FGF-8a is often used for the differentiation of induced [pluripotent](#) stem cells, embryonic stem cells, and neural stem cells. FGF-8a is a spliced form of FGF-8, a heparin-binding protein that targets mammary and non-mammary cells expressing the FGF receptors.

Qkine mouse/rat FGF-8a is a 21.3 kDa highly pure, bioactive recombinant protein produced in an animal-free expression system. This protein is [animal-origin free](#) (AOF), carrier protein-free, tag-free, and non-glycosylated to ensure a pure homogenous protein with exceptional lot-to-lot consistency.

Alternative protein names

AIGF, AIGFKAL6, Androgen-induced growth factor, FGF8, FGF-8, Fibroblast Growth Factor - 8, HBGF-8, Heparin Binding Growth Factor - 8, MGC149376, FGF8a, FGF 8a, Qk20

Molecular weight

21.3 kDa

Protein Uniprot number

High purity FGF-8a protein (Uniprot: P37237)

Species reactivity

- mouse
- rat
- species similarity:
- human - 100%
- bovine - 100%
- porcine - 100%

Product Information

- >98%, by SDS-PAGE quantitative densitometry
- Expressed in *E. coli*
- Animal origin-free (AOF) and carrier protein-free
- Manufactured in our Cambridge, UK laboratories
- Lyophilized from HEPES, NaCl, mannitol

Reconstitution instructions

- Resuspend in sterile-filtered water at >50 µg/ml

Featured applications

- Differentiation of midbrain dopaminergic neurons
- Differentiation of iPSC derived neural progenitors to neurons


Further quality assays

- Mass spectrometry: single species with expected mass
- Recovery from stock vial: >95%
- Endotoxin: <0.05 EU/µg protein

Scientific Information


Bioactivity

FGF-8a activity was determined using the Promega serum response element luciferase reporter assay in HEK293T cells. Cells were treated in triplicate with a serial dilution of FGF-8a in the presence of 10 µg/ml heparin for 3 hours. Firefly luciferase activity was measured and normalized to the control Renilla luciferase activity. Data from lot #104406. EC50 = 110 ng/ml (5.2 nM).



Purity

FGF-8a migrates as a single band at 21.3 kDa in non-reducing (NR) conditions and upon reduction (R). Purified recombinant protein (3 µg) was resolved using 15% w/v SDS-PAGE in reduced (+β-mercaptoethanol, R) and non-reduced (NR) conditions and stained with Coomassie Brilliant Blue R250. Data from lot #104412.



Original product page: <https://ryan.calliope-alpha.ts.net/product/recombinant-mouse-rat-fgf-8a-protein-qk020/>

PDF generated: 12 May 2026

Copyright © 2026 by Qkine Ltd. All rights reserved including graphics and images.